


- 25 Benetos A, Rudnichi A, Safar M, *et al*. Pulse pressure and cardiovascular mortality in normotensive and hypertensive subjects. *Hypertension* 1998;**32**:560–4.
- 26 van Trijp MJ, Grobbee DE, Peeters PH, *et al*. Average blood pressure and cardiovascular disease-related mortality in middle-aged women. *Am J Hypertens* 2005;**18**:197–201.
- 27 Schram MT, Chaturvedi N, Fuller JH, *et al*. Pulse pressure is associated with age and cardiovascular disease in type 1 diabetes: the Eurodiab Prospective Complications Study. *J Hypertens* 2003;**21**:2035–44.
- 28 Park MK, Menard SM. Accuracy of blood pressure measurement by the Dinamap monitor in infants and children. *Pediatrics* 1987;**79**:907–14.
- 29 Colan SD, Fujii A, Borow KM, *et al*. Noninvasive determination of systolic, diastolic and end-systolic blood pressure in neonates, infants and young children: comparison with central aortic pressure measurements. *Am J Cardiol* 1983;**52**:867–70.
- 30 O'Brien E. Demise of the mercury sphygmomanometer and the dawning of a new era in blood pressure measurement. *Blood Press Monit* 2003;**8**:19–21.
- 31 O'Brien E, Petrie J, Littler W, *et al*. The British Hypertension Society protocol for the evaluation of automated and semi-automated blood pressure measuring devices with special reference to ambulatory systems. *J Hypertens* 1990;**8**:607–19.
- 32 O'Brien E, Mee F, Atkins N, *et al*. Short report: accuracy of the Dinamap portable monitor, model 8100 determined by the British Hypertension Society protocol. *J Hypertens* 1993;**11**:761–3.
- 33 Wattigney WA, Webber LS, Lawrence MD, *et al*. Utility of an automatic instrument for blood pressure measurement in children. The Bogalusa Heart Study. *Am J Hypertens* 1996;**9**:256–62.
- 34 Barker ME, Shiell AW, Law CM. Evaluation of the Dinamap 8100 and Omron M1 blood pressure monitors for use in children. *Paediatr Perinat Epidemiol* 2000;**14**:179–86.
- 35 Park MK, Menard SW, Yuan C. Comparison of auscultatory and oscillometric blood pressures. *Arch Pediatr Adolesc Med* 2001;**155**:50–3.
- 36 Jin RZ, Donaghue KC, Fairchild J, *et al*. Comparison of Dinamap 8100 with sphygmomanometer blood pressure measurement in a prepubertal diabetes cohort. *J Paediatr Child Health* 2001;**37**:545–9.
- 37 O'Brien E, Atkins N. Inaccuracy of the Dinamap 8100 portable monitor. *Lancet* 1997;**349**:1026.
- 38 Friedman B. Accuracy of Dinamap monitors. *Lancet* 1997;**350**:217–18.
- 39 National Blood Pressure Education Working Group on High Blood Pressure in Children and Adolescents. Fourth report on the diagnosis, evaluation and treatment of high blood pressure in children and adolescents: a working group report from the National High Blood Pressure Education Program. *Pediatrics* 2004;**114**:555–76.
- 40 St George IM, Williams SM, Silva PA. The stability of high blood pressure in Dunedin children: an eight year longitudinal study. *NZ Med J* 1990;**103**:115–17.
- 41 de Swiet M, Foyers P, Shinebourne EA. Blood pressure in first 10 years of life: the Brompton study. *BMJ* 1992;**304**:23–6.
- 42 Chinn S, Rona RJ. Prevalence and trends in overweight and obesity in three cross sectional studies of British children 1974–94. *BMJ* 2001;**322**:24–6.

## IMAGES IN PAEDIATRICS.....

doi: 10.1136/adc.2006.112771

### An interesting facial rash



**Figure 1** Informed consent was obtained for publication of this figure.

**A** 7-year-old boy presented to paediatrics with a 3-week history of a worsening facial rash, which was intermittently itchy. He was otherwise well. He had two guinea pigs as pets. On examination, an extensive scaling erythema was noticed with a definite edge involving the upper eyelids, the bridge of the nose and extending onto both cheeks (see fig 1).

A provisional diagnosis of tinea faciei was made; however, cutaneous lupus was also considered. While mycology results were awaited, topical terbinafine was given, with little effect. Microscopy revealed a dermatophyte infection with *Trichophyton mentagrophytes*, and a 3-week course of oral terbinafine (125 mg oral dosage once daily) was given. The rash resolved completely, leaving post-inflammatory hyperpigmentation only.

Tinea facialis/faciei is a dermatophytosis of the glabrous facial skin, characterised by a well-circumscribed, often asymmetric, erythematous patch with an elevated border and central regression. It may be asymptomatic or present with pruritus, or, occasionally, photosensitivity that may lead to diagnostic confusion with cutaneous lupus.<sup>1</sup> It is the most commonly misdiagnosed dermatophytosis. Other differential diagnoses include eczema, seborrhoeic dermatitis<sup>2</sup> and rosacea.

It is most common in children, with predisposing factors including exposure to animals, chronic topical steroid use and spread from tinea capitis. The most frequent organisms involved are *T mentagrophytes*, *T rubrum* and *T tonsurans*. However, cases caused by *Microsporum audouinii* and *M canis* occur worldwide. Most cases are given short-term oral antifungal treatment, but milder cases may respond to topical imidazoles. Affected animals and family members should also be treated.

**Caroline A Love**

Department of Dermatology, Rowan House, Whiston Hospital, Merseyside, UK

**John A Sills, Judith M Ellison**

St Helens and Knowsley NHS Trust, Merseyside, UK

Correspondence to: Dr C A Love, Department of Dermatology, Rowan House, Whiston Hospital, Warrington Road, Prescott, Merseyside L35 5DR, UK; drcalove@hotmail.com

Competing interests: None declared.

## REFERENCES

- 1 Cirillo-Hyland V, Humphreys T, Elenitsas R. Tinea faciei. *J Am Acad Dermatol* 1993;**29**:119–20.
- 2 Gorani A, Oriani A, Cambiaghi S. Seborrhoeic dermatitis-like tinea faciei. *Pediatr Dermatol* 2005;**22**:243–4.